


# Polygone de longueur minimal dynamique

J.-O. Lachaud, X. Provençal




23 juillet 2020


# Minimum Length Polygon




# Minimum Length Polygon




# Minimum Length Polygon



# Minimum Length Polygon




# Minimum Length Polygon



## Definition

Given a digital contour  $C$ , its *inner* (resp. *outer*) contour  $IC(C)$  (resp.  $OC(C)$ ) is the erosion (resp. dilatation) of the body of  $I(C)$  by the open unit square centred on  $(0,0)$ .


# Minimum Length Polygon



## Definition

Given a digital contour  $C$ , its *inner* (resp. *outer*) contour  $IC(C)$  (resp.  $OC(C)$ ) is the erosion (resp. dilatation) of the body of  $I(C)$  by the open unit square centred on  $(0,0)$ .

# Minimum Length Polygon




## Definition

Given a digital contour  $C$ , its *inner* (resp. *outer*) contour  $IC(C)$  (resp.  $OC(C)$ ) is the erosion (resp. dilatation) of the body of  $I(C)$  by the open unit square centred on  $(0,0)$ .



# Minimum Length Polygon





## Definition

The *minimum length polygon* of  $C$  is a subset  $P \in \mathbb{R}^2$  such that,

$$P = \arg \min_{A \in \mathcal{A}, IC(C) \subseteq A, \partial A \subseteq OC(C) \setminus IC(C)^\circ} \text{Per}(A)$$



where  $\mathcal{A}$  is the family of simply connected compact sets of  $\mathbb{R}^2$ .

# Minimum Length Polygon



The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :



# Minimum Length Polygon



The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :

- unique ;

# Minimum Length Polygon





The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :

- unique ;
- a good length estimator<sup>1</sup> ;

<sup>1</sup> Proved to be convergent on convex shapes.

# Minimum Length Polygon





The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :

- unique ;
- a good length estimator<sup>1</sup> ;
- a good tangent estimator ;

<sup>1</sup> Proved to be convergent on convex shapes.

# Minimum Length Polygon





The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :

- unique ;
- a good length estimator<sup>1</sup> ;
- a good tangent estimator ;
- characteristic of the shape's convexity ;

<sup>1</sup> Proved to be convergent on convex shapes.

# Minimum Length Polygon



The MLP is a polygonal line whose vertices are centers of pixels along the inner or the outer contour, also :

- unique ;
- a good length estimator<sup>1</sup> ;
- a good tangent estimator ;
- characteristic of the shape's convexity ;
- reversible<sup>2</sup>.

<sup>1</sup> Proved to be convergent on convex shapes.

<sup>2</sup> If aligned vertices are considered.

MLP is computable in time linear with respect of the length of  $C$ .

- J.-O. Lachaud, X. Provençal, *Two linear-time algorithms for computing the minimum length polygon of a digital contour*, Discrete Applied Mathematics (DAM), 2011.



# Segmentation using deformable models









Fig. 4. Example of the minimization process using the Greedy1 algorithm. The gradient is computed with the Canny-Deriche method with scale coefficient 0.2. The input image represents a half-plane. (First row) Initialisation of the DDM. (Second row) Results of the minimisation process, the  $\alpha$  coefficient used is equal to 0. (Third row) Results with  $\alpha = 200$ . (Fifth row) Results with  $\alpha = 300$ .


- F. de Veilleville and J.-O. Lachaud, *Digital Deformable Model Simulating Active Contours*, in proc. DGCI2009, LNCS 5810, p.203-216, 2009.

# Segmentation using deformable models




- G. Damiand, A. Dupas and J.-O. Lachaud, *Combining Topological Maps, Multi-Label Simple Points, and Minimum-Length Polygons for Efficient Digital Partition Model*, in proc. IWCMIA2011, LNCS 6636, p. 208-221, 2011.


# Flip a pixel




# Flip a pixel




# Flip a pixel




# Flip a pixel



# Flip a pixel



# Flip a pixel





# Reversible polygonal representation

Goal : represent a digital contour  $C$  using a polygon whose vertices are centers of pixels either on the inner contour  $IC(C)$  or on the outer contour  $OC(C)$ .

# Reversible polygonal representation

Goal : represent a digital contour  $C$  using a polygon whose vertices are centers of pixels either on the inner contour  $IC(C)$  or on the outer contour  $OC(C)$ .

## Definition

A *grid-vector* is a triplet  $x = ((p, q), k, \delta) \in \mathbb{N}^2 \times \mathbb{N} \times \mathbb{B}$ . where

- $\gcd(p, q) = 1$ ,  $q/p$  is the *slope* of  $x$  (with  $1/0 = \infty$ ),
- $k \geq 1$  is its number of repetitions
- the boolean  $\delta$  indicates if  $x$  has one endpoint on the inner contour and one on the outer.

Notation :  $((p, q), k, \delta) = \begin{cases} (p, q)^k & \text{if } \delta \text{ is false,} \\ \widetilde{(p, q)}^k & \text{otherwise.} \end{cases}$

# Reversible polygonal representation

Geometric interpretation of grid-vectors.


## Definition

A *context* is an ordered pair of letters  $(a, b)$  among  $\{(0, 1), (1, 2), (2, 3), (3, 0), (0, 3), (3, 2), (2, 1), (1, 0)\}$ .

Given a context  $(a, b)$ , a grid-vectors defines the following vector as follow :

$$\overrightarrow{(p, q)}^{(a, b)} = k(p \vec{a} + q \vec{b}),$$

$$\overleftarrow{(p, q)}^{(a, b)} = k(p \vec{b} + q \vec{a}).$$



# Reversible polygonal representation

Geometric interpretation of grid-vectors.



## Definition

A *context* is an ordered pair of letters  $(a, b)$  among  $\{(0, 1), (1, 2), (2, 3), (3, 0), (0, 3), (3, 2), (2, 1), (1, 0)\}$ .


Given a context  $(a, b)$ , a grid-vectors defines the following vector as follow :

$$\overrightarrow{(p, q)}^{(a, b)} = k(p \vec{a} + q \vec{b}),$$


$$\overleftarrow{(p, q)}^{(a, b)} = k(p \vec{b} + q \vec{a}).$$




$(3, 2)^1$



$(2, 3)^1$



$(3, 2)^1$



$(3, 2)^1$



# Reversible polygonal representation

Notations :

- $\overset{(a,b)}{\overrightarrow{\sigma}^-} = \overset{(a,b)}{\overrightarrow{\sigma}^+} = (0,0)$ .


- Let  $x = ((p, q), k, \delta)$ ,  $x(a, b) = \begin{cases} (b, a) & \text{if } \delta \text{ is true,} \\ (a, b) & \text{otherwise.} \end{cases}$

# Reversible polygonal representation

Notations :

- $\overrightarrow{\sigma^-} = \overrightarrow{\sigma^+} = (0, 0)$ .

- Let  $x = ((p, q), k, \delta)$ ,  $x(a, b) = \begin{cases} (b, a) & \text{if } \delta \text{ is true,} \\ (a, b) & \text{otherwise.} \end{cases}$





$[(2, 3), (3, 1), \widetilde{(1, 1)}, (2, 3), (3, 1)]$

# Reversible polygonal representation

Notations :

- $\overrightarrow{\sigma^-} = \overrightarrow{\sigma^+} = (0, 0)$ .

- Let  $x = ((p, q), k, \delta)$ ,  $x(a, b) = \begin{cases} (b, a) & \text{if } \delta \text{ is true,} \\ (a, b) & \text{otherwise.} \end{cases}$



$[(2, 3), (3, 1), \widetilde{(1, 1)}, (2, 3), (3, 1)]$



# Reversible polygonal representation

Notations :

- $\overset{(a,b)}{\rightarrow}{\sigma}^- = \overset{(a,b)}{\rightarrow}{\sigma}^+ = (0,0)$ .

- Let  $x = ((p, q), k, \delta)$ ,  $x(a, b) = \begin{cases} (b, a) & \text{if } \delta \text{ is true,} \\ (a, b) & \text{otherwise.} \end{cases}$

From grid-curves to polygons.


A grid-curve  $\Gamma = [l_0, l_1, \dots, l_{n-1}]$ , a context  $(a_0, b_0)$  and a start point  $P_0$  define a polygonal curve  $P_\Gamma = [P_0, P_1, \dots, P_n]$  as follow :

$$P_{i+1} = P_i + \overset{(a_i, b_i)}{\rightarrow}{l_i} \quad \text{and} \quad (a_{i+1}, b_{i+1}) = l_i(a_i, b_i).$$

By fixing the first point on the inside or outside polygon, a discrete contour is defined unambiguously.


# Reversible polygonal representation

Some words about Christoffel words.




# Reversible polygonal representation

Some words about Christoffel words.




# Reversible polygonal representation

Some words about Christoffel words.



The *standard factorization* of a Christoffel word  $w$  is the only factorization  $w = uv$  where  $u$  and  $v$  are both Christoffel words.

# Reversible polygonal representation



$$\begin{aligned} \alpha &= [z_0; z_1, z_2, \dots] \\ &= z_0 + \frac{1}{z_1 + \frac{1}{z_2 + \frac{1}{z_3 + \dots}}} \end{aligned}$$

The Christoffel word  $c_n$  of slope  $[z_0; z_1, z_2, \dots, z_n]$  is given recursively by :

$$c_n = \begin{cases} c_{2m-2} c_{2m-1}^{z_{2m}} & \text{if } n = 2m, \\ c_{2m}^{z_{2m+1}} c_{2m-1} & \text{if } n = 2m + 1. \end{cases}$$

where  $c_{-1} = 2$ , and  $c_{-2} = 1$ ,

# Reversible polygonal representation


Let  $C_{q/p}^{(a,b)}$  be the Christoffel word of slope  $q/p$  over the alphabet  $(a, b)$ .

A grid-curve  $\Gamma = [l_0, l_1, \dots, l_{n-1}]$  defines a digital contour by gluing all  $F_{(a_i, b_i)}(l_i)$  defined as follow :

$$\begin{aligned} F_{(a,b)}((p, q)^k) &= \left(C_{q/p}^{(a,b)}\right)^k, & F_{(a,b)}(\sigma^-) &= \bar{b}, \\ F_{(a,b)}(\widetilde{(p, q)^k}) &= a\bar{b}F_{(b,a)}((p, q)^k), & F_{(a,b)}(\sigma^+) &= a. \end{aligned}$$


The interpixel path  $F(\Gamma)$  is obtained from by removing *cancellations* that are factors of the form  $a\bar{a}$ .

# Not unique




$$\Gamma = [(3, 2), \sigma^+, (3, 2)],$$

$$\begin{aligned} F_{(0,1)}(\Gamma) &= F_{(0,1)}((3, 2)) \cdot F_{(0,1)}(\sigma^+) \cdot F_{(3,0)}((3, 2)) \\ &= 00101 \cdot 0 \cdot 33030 \end{aligned}$$



$$\Gamma = [(3, 2), \sigma^-, (3, 2)],$$


$$\begin{aligned} F_{(0,1)}(\Gamma) &= F_{(0,1)}((3, 2)) \cdot F_{(0,1)}(\sigma^-) \cdot F_{(1,2)}((3, 2)) \\ &= 00101 \cdot 3 \cdot 11212 \end{aligned}$$



$$\Gamma = [\widetilde{(3, 2)}, (3, 2)],$$

$$\begin{aligned} F_{(0,1)}(\Gamma) &= F_{(0,1)}(\widetilde{(3, 2)}) \cdot F_{(1,0)}((3, 2)) \\ &= 03 \cdot 11010 \cdot 11010 \end{aligned}$$

# Reversible polygonal representation




$$\begin{aligned}
 F_{(0,1)} \left( ((1, 2), 2, \text{false}) \right) &= (011)^2, \\
 F_{(0,1)} \left( ((1, 2), 1, \text{true}) \right) &= 03 \cdot 100, \\
 F_{(1,0)} \left( ((3, 1), 1, \text{true}) \right) &= 12 \cdot 0001, \\
 F_{(0,1)} \left( ((1, 0), 2, \text{false}) \right) &= 0^2, \\
 F_{(0,1)} (\sigma^+) &= 0, \\
 F_{(3,0)} \left( ((3, 2), 1, \text{true}) \right) &= 32 \cdot 00303, \\
 F_{(0,3)} (\sigma^-) &= 1, \\
 F_{(3,2)} \left( ((2, 1), 1, \text{false}) \right) &= 332, \\
 F_{(3,2)} \left( ((2, 1), 1, \text{true}) \right) &= 30 \cdot 223,
 \end{aligned}$$

$$\begin{aligned}
 w &= 011011 \cdot 0\cancel{31}00 \cdot 1\cancel{20}001 \cdot 00 \cdot 0 \cdot 3\cancel{20}030\cancel{31}332 \cdot 3\cancel{02}23, \\
 &= 011011 \cdot 000 \cdot 1001 \cdot 00 \cdot 0 \cdot 3030 \cdot 3 \cdot 32 \cdot 323.
 \end{aligned}$$



# Reversible polygonal representation



$$\begin{aligned}
 F_{(0,1)} \left( ((1, 2), 2, \text{false}) \right) &= (011)^2, \\
 F_{(0,1)} \left( ((1, 2), 1, \text{true}) \right) &= 03 \cdot 100, \\
 F_{(1,0)} \left( ((3, 1), 1, \text{true}) \right) &= 12 \cdot 0001, \\
 F_{(0,1)} \left( ((1, 0), 2, \text{false}) \right) &= 0^2, \\
 F_{(0,1)} (\sigma^+) &= 0, \\
 F_{(3,0)} \left( ((3, 2), 1, \text{true}) \right) &= 32 \cdot 00303, \\
 F_{(0,3)} (\sigma^-) &= 1, \\
 F_{(3,2)} \left( ((2, 1), 1, \text{false}) \right) &= 332, \\
 F_{(3,2)} \left( ((2, 1), 1, \text{true}) \right) &= 30 \cdot 223,
 \end{aligned}$$

$$\begin{aligned}
 w &= 011011 \cdot 0 \cancel{31} 00 \cdot 1 \cancel{20} 001 \cdot 00 \cdot 0 \cdot 3 \cancel{20} 030 \cancel{31} 332 \cdot 3 \cancel{02} 23, \\
 &= 011011 \cdot 000 \cdot 1001 \cdot 00 \cdot 0 \cdot 3030 \cdot 3 \cdot 32 \cdot 323.
 \end{aligned}$$

## Definition

Two grid-curves  $\Gamma$  and  $\Gamma'$  are *equivalent*, if they define the same digital contour and ends in the same orientation.

The MLP of the digital contour  $C$  is the shortest grid-curve in the equivalence class defined by  $C$ .






# Relative orientation of grid-segements

## Notation

Given  $x = ((p, q), k, \delta_x)$  and  $y = ((r, s), l, \delta_y)$ ,

$$x \otimes y = \begin{cases} ps - qr & \text{if } \delta_y \text{ is false,} \\ pr - qs & \text{if } \delta_y \text{ is true.} \end{cases}$$

### Three cases

$x \otimes y = 0$	$x \otimes y < 0$	$x \otimes y > 0$
		
$[(3, 2), (3, 2)]$	$[(2, 3), (2, 1)]$	$[(3, 1), (2, 3)]$
		
	$[\widetilde{(1, 3)}, \widetilde{(2, 3)}]$	$[\widetilde{(3, 2)}, \widetilde{(2, 1)}]$




# Relative orientation of grid-segements

## Notation

Given  $x = ((p, q), k, \delta_x)$  and  $y = ((r, s), l, \delta_y)$ ,

$$x \otimes y = \begin{cases} ps - qr & \text{if } \delta_y \text{ is false,} \\ pr - qs & \text{if } \delta_y \text{ is true.} \end{cases}$$

### Three cases

$x \otimes y = 0$	$x \otimes y < 0$	$x \otimes y > 0$
		
$[(3, 2)^2]$	$[(2, 3), (2, 1)]$ $[\widetilde{(1, 3)}, \widetilde{(2, 3)}]$	$[(3, 1), (2, 3)]$ $[\widetilde{(3, 2)}, \widetilde{(2, 1)}]$

# Merge case : $x \otimes y = 1$

Let  $x = ((p, q), k, \delta_x)$  and  $y = ((r, s), l, \delta_y)$  with  
 $\delta_y = \text{false}$  and  $\min(k, l) = 1$


$$x \otimes y = 1,$$

or


$$\delta_y = \text{true} \text{ and } l = 1$$

then


$$[x, y] \equiv [z] \text{ where } z = \begin{cases} ((kp + lr, kq + ls), 1, \delta_x) & \text{if } \delta_y = \text{false.} \\ ((kp + ls, kq + lr), 1, \neg\delta_x) & \text{otherwise.} \end{cases}$$




$$[(2, 1)^2, (1, 1)]$$




$$[(2, 1), (1, 1)^2]$$




$$[(1, 1)^2, \widetilde{(2, 1)}]$$




$$[\widetilde{(2, 1)^2}, (1, 1)]$$




$$[(5, 3)]$$



$$[(4, 3)]$$




$$[\widetilde{(4, 3)}]$$



$$[\widetilde{(5, 3)}]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (\widetilde{3, 1}), (2, 1)^3]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2), (\widetilde{1, 0}), (2, 1)^3]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2), (\widetilde{1, 0}), (2, 1), (2, 1)^2]$$




$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2), (\widetilde{1, 0}), (\widetilde{1, 1}), (\widetilde{1, 0}), (2, 1)^2]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2), (\widetilde{1, 0}), (\widetilde{1, 0}), (0, 1), (\widetilde{1, 0}), (2, 1)^2]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2), (1, 1)(0, 1), (\widetilde{5, 2})]$$

# Split and merge case : $x \otimes y > 1$


$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (1, 2)^2, (\widetilde{5, 2})]$$

# Split and merge case : $x \otimes y > 1$

$$(8, 3) \otimes (2, 1)^3 = 2.$$



$$[(8, 3), (2, 1)^3] \equiv [(\widetilde{2, 5}), (\widetilde{9, 4})]$$

# How to split ?

## Notation

Let  $x = ((p, q), 1, \text{false})$  and  $q/p = [u_0; u_1, \dots, u_n]$ .

- $q_i/p_i = [u_0; u_1, \dots, u_i]$ ,      •  $x_{-1} = ((0, 1), 1, \text{false})$ ,
- $x_i = ((p_i, q_i), 1, \text{false})$ ,      •  $x_{-2} = ((1, 0), 1, \text{false})$ .

## Definition


The *basic splitting* of the grid-vector  $x_n$  is the grid-curve :

$$s(x_n) = \begin{cases} [x_{2m-2}, x_{2m-1}^{u_{2m}}] & \text{if } n = 2m, \\ [x_{2m}^{u_{2m+1}}, x_{2m-1}] & \text{if } n = 2m+1, \end{cases}$$


A grid-vector and its basic splittings both define the same interpixel path.

$$s(x) = [y, z] \implies y \otimes z = 1.$$

# How to split?




# How to split ?






# How to split?



$$5/7 = [0; 1, 2, 2],$$


$$2/3 = [0; 1, 2],$$

$$1/1 = [0; 1]$$



$$[(7, 5)] \equiv [(3, 2)^2, (1, 1)]$$


# How to split?



$$5/7 = [0; 1, 2, 2],$$


$$2/3 = [0; 1, 2],$$

$$1/1 = [0; 1]$$




$$[(7, 5)] \equiv [(3, 2)^2, (1, 1)] \equiv [(\widetilde{1, 1}), (2, 3), (\widetilde{3, 2})]$$

# Flip a pixel




# Flip a pixel



$$[(\widetilde{3, 8}), (\widetilde{2, 1})^3] \equiv [(\widetilde{1, 2}), (1, 3), (1, 3), (\widetilde{2, 1})^3]$$

- 1 Split grid-segments until one ends exactly on the pixel to flip. Let  $x = ((p, q), 1, \delta_x)$  be the grid segment right before and  $y = ((r, s), 1, \delta_y)$  be the grid-vector right after.

# Flip a pixel




$$[(\widetilde{3, 8}), (\widetilde{2, 1})^3] \equiv [(\widetilde{1, 2}), (1, 3), (1, 3), (\widetilde{2, 1})^3]$$

$$\neq [(\widetilde{1, 2}), (\widetilde{3, 1}), (\widetilde{1, 3}), (\widetilde{2, 1})^3]$$

- 1 Split grid-segments until one ends exactly on the pixel to flip. Let  $x = ((p, q), 1, \delta_x)$  be the grid segment right before and  $y = ((r, s), 1, \delta_y)$  be the grid-vector right after.
- 2 Replace  $x$  by  $((q, p), 1, -\delta_x)$ .
- 3 Replace  $y$  by  $((r, s), 1, -\delta_y)$ .

# Flip a pixel




$$[(\widetilde{3, 8}), (\widetilde{2, 1})^3] \equiv [(\widetilde{1, 2}), (1, 3), (1, 3), (\widetilde{2, 1})^3]$$

$$\neq [(\widetilde{1, 2}), (\widetilde{3, 1}), (\widetilde{1, 3}), (\widetilde{2, 1})^3] \equiv [(5, 2), (\widetilde{1, 3}), (\widetilde{2, 1})^3]$$

- 1 Split grid-segments until one ends exactly on the pixel to flip. Let  $x = ((p, q), 1, \delta_x)$  be the grid segment right before and  $y = ((r, s), 1, \delta_y)$  be the grid-vector right after.
- 2 Replace  $x$  by  $((q, p), 1, \neg\delta_x)$ .
- 3 Replace  $y$  by  $((r, s), 1, \neg\delta_y)$ .

# Flip a pixel




$$[(\widetilde{3, 8}), (\widetilde{2, 1})^3] \equiv [(\widetilde{1, 2}), (1, 3), (1, 3), (\widetilde{2, 1})^3]$$

$$\neq [(\widetilde{1, 2}), (\widetilde{3, 1}), (\widetilde{1, 3}), (\widetilde{2, 1})^3] \equiv [(5, 2), (\widetilde{1, 3}), (\widetilde{2, 1})^3]$$

- 1 Split grid-segments until one ends exactly on the pixel to flip. Let  $x = ((p, q), 1, \delta_x)$  be the grid segment right before and  $y = ((r, s), 1, \delta_y)$  be the grid-vector right after.
- 2 Replace  $x$  by  $((q, p), 1, \neg\delta_x)$ .
- 3 Replace  $y$  by  $((r, s), 1, \neg\delta_y)$ .


# Flip a pixel on a flat part



$[(2, 1), (1, 0)^6, \sigma^+, (1, 2)]$




# Flip a pixel on a flat part




$[(2, 1), (1, 0)^3, (1, 0)^3, \sigma^+, (1, 2)]$

# Flip a pixel on a flat part



$$[(2, 1), (1, 0)^2, \widetilde{(0, 1)}, \widetilde{(1, 0)}^3, \sigma^+, (1, 2)]$$

# Flip a pixel on a flat part




$[(2, 1), (1, 0)^3, (1, 0)^3, \sigma^+, (1, 2)]$





# Flip a pixel on a flat part




$$[(2, 1), (1, 0)^3, \underbrace{\sigma^-, (1, 0), \sigma^+, \sigma^+, (1, 0), \sigma^-}_{\text{bumb}}, (1, 0)^3, \sigma^+, (1, 2)]$$

How to simplify  $\sigma^-$  ?

- Cancellation :  $[\sigma^-, \sigma^+] \equiv [\sigma^+, \sigma^-] \equiv []$

# Flip a pixel on a flat part



$$[(2, 1), (1, 0)^3, \underbrace{\sigma^-, (1, 0), \sigma^+, \sigma^+, (1, 0), \sigma^-}_{\text{bumb}}, (1, 0)^3, \sigma^+, (1, 2)]$$

How to simplify  $\sigma^-$  ?

- Cancellation :  $[\sigma^-, \sigma^+] \equiv [\sigma^+, \sigma^-] \equiv []$
- Split the grid-edges in order to have a local part build only with  $\{\sigma^+, \sigma^-, (1, 0), (0, 1), \widetilde{(1, 0)}, \widetilde{(0, 1)}\}$ . Operators  $\sigma^-$  are then simplify using local rules such as :

$$[(1, 0), \sigma^-, (1, 0), \sigma^+] \equiv [(1, 1)] \text{ and } [\sigma^-, (1, 0)^k, \sigma^+] \equiv [(0, 1)^k]$$

## Proposition

A grid-curve defining a digital contour may be simplified to a MLP using local rules.



## Proposition

A grid-curve defining a digital contour may be simplified to a MLP using local rules.

## Proposition

Given a grid-curve that is the MLP of a digital contour, this contour may be modified by adding or removing one pixel and its MLP updated in time sub-linear with respect to the length of the modified part of the MLP.

Implemente in project *ImaGene* available at  
[gforge.liris.cnrs.fr/projects/imagene](http://gforge.liris.cnrs.fr/projects/imagene)

*MERCI !*